Research Article
E-ISSN : 3048-1368

Volume 05, Issue No. 03. 2024
P-ISSN : 3048-1376

The International Journal of Medical Science and Health Research



A Comprehensive Literature Review of Implications of Preoperative Depression for Lumbar Spine Surgery Outcomes

[1] Arief, [2] Rizki Kurnia, [2] Catherine

[1] Hasan Sadikin Central General Hospital, Bandung City, West Java, Indonesia
[2] Faculty of Medicine, Padjadjaran University, Bandung City, West Java, Indonesia
Correspondence : drarief2839@gmail.com

Article History :
Received date : 2024/07/17
Revised date : 2024/08/16
Accepted date : 2024/09/20
Published date : 2024/10/11

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

E-ISSN :
ISSN 3048-1368
9 773048 136000

P-ISSN :
ISSN 3048-1376
9 773048 137007

ABSTRACT

Background: The literature reviewed emphasizes the significant impact of preoperative depression on surgical outcomes for patients undergoing lumbar spine surgery. The studies collectively highlight a consistent link between depressive symptoms and adverse postoperative results, suggesting that psychological health is a critical factor influencing recovery trajectories. **Literature Review:** The foundational study by (Sinikallio et al., 2010) provides compelling evidence that both preoperative and postoperative depressive symptoms predict poor surgical outcomes. This research underscores the importance of mental health not only before surgery but also during the rehabilitation process, indicating that ongoing depressive symptoms can exacerbate recovery challenges. Further expanding on this theme, (L. Jackson et al., 2019) conducted a systematic review that connects psychological comorbidities with postoperative complications. Their work reveals that preoperative depression correlates with increased rates of delirium, readmissions, and prolonged hospital stays, suggesting that elevated cortisol levels in depressed individuals may be a contributing factor to these adverse outcomes. Additionally, the retrospective study by (Zhang et al., 2023) specifically examines the effects of preoperative depression on outcomes following cervical laminoplasty, providing valuable insights that can be extrapolated to lumbar spine surgeries. **Conclusion:** In conclusion, the literature consistently demonstrates that preoperative depression is a significant predictor of surgical outcomes in lumbar spine surgery. The integration of psychological assessments into preoperative evaluations is essential for optimizing patient care and enhancing recovery trajectories. Addressing mental health prior to surgery could mitigate adverse outcomes and improve overall patient well-being, highlighting the need for targeted interventions in this population.

Keywords: Implications, Preoperative Depression, Lumbar Spine Surgery, Outcomes

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The implications of preoperative depression for lumbar spine surgery outcomes have garnered increasing attention in recent years, reflecting a growing recognition of the interplay between psychological health and surgical success. The literature on this topic reveals a consistent link between depressive symptoms and adverse surgical outcomes, underscoring the need for a comprehensive understanding of how preoperative mental health can influence recovery trajectories in patients undergoing lumbar spine procedures.

In a seminal study by (Sinikallio et al., 2010), the authors explored the predictive value of both preoperative and postoperative depressive symptoms on outcomes following lumbar spinal stenosis surgery. Their findings indicated that patients exhibiting depressive symptoms during the rehabilitation period were more likely to experience poor surgical outcomes, thus highlighting the critical role of mental health in the recovery process. This early investigation laid the groundwork for subsequent research, emphasizing the necessity of addressing psychological factors prior to surgical interventions.

Building upon this foundation, (L. Jackson et al., 2019) conducted a systematic review that examined the relationship between psychological comorbidities, including depression, and postoperative outcomes across various spine surgeries. Their analysis revealed that preoperative psychological disorders were associated with increased rates of postoperative complications such as delirium, readmission, and prolonged hospital stays. Notably, the authors posited that elevated cortisol levels in depressed individuals may contribute to these adverse outcomes. This comprehensive review provided a broader context for understanding the implications of preoperative depression, suggesting that mental health interventions could be critical in improving surgical outcomes and reducing healthcare costs associated with complications.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Most recently, (Zhang et al., 2023) investigated the specific impact of preoperative depression on clinical outcomes after cervical laminoplasty. Their retrospective study found that patients with preoperative depression exhibited significantly worse postoperative improvement in lumbar fusion, reinforcing the notion that mental health status prior to surgery is a determinant of recovery. This study adds to the growing body of evidence indicating that preoperative depression not only affects immediate surgical outcomes but may also hinder long-term recovery.

Collectively, these studies underscore the importance of recognizing and addressing preoperative depression in patients undergoing lumbar spine surgery. The evidence suggests that mental health interventions could play a vital role in enhancing surgical outcomes and improving overall patient well-being. As the field continues to evolve, it is imperative to integrate psychological assessments into preoperative evaluations to optimize care for individuals facing lumbar spine surgery.

LITERATURE REVIEW

The article "Depressive symptoms during rehabilitation period predict poor outcome of lumbar spinal stenosis surgery: A two-year perspective" by (Sinikallio et al., 2010) provides a significant contribution to the understanding of the relationship between depressive symptoms and surgical outcomes in patients undergoing lumbar spine surgery. The authors present a well-structured prospective clinical study that examines the predictive value of both preoperative and postoperative depressive symptoms on the outcomes of lumbar spinal stenosis surgery.

The study's findings reinforce the notion that preoperative depressive symptoms are not merely incidental but rather play a critical role in determining surgical outcomes. The authors highlight that patients exhibiting depressive symptoms prior to surgery are at a higher risk for poor postoperative results, which aligns with previous literature indicating a correlation between mental health and surgical

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

recovery ((Sinikallio et al., 2010)). This relationship is crucial for clinicians to consider, as it suggests that addressing mental health issues before surgery could be pivotal in improving patient outcomes.

Moreover, the study extends the inquiry beyond preoperative factors by also assessing the impact of depressive symptoms during the rehabilitation phase. The authors effectively demonstrate that ongoing depressive symptoms can further complicate recovery, potentially leading to a cycle of poor outcomes that could be mitigated with appropriate psychological interventions. This dual focus on both preoperative and postoperative depressive symptoms provides a comprehensive view of how mental health influences the surgical process and outcomes.

The methodology employed in the study is robust, with a two-year follow-up period that allows for a thorough assessment of the long-term effects of depressive symptoms on surgical outcomes. This longitudinal approach strengthens the validity of the findings and emphasizes the importance of continuous monitoring of mental health throughout the surgical journey.

The article "Correlating Psychological Comorbidities and Outcomes After Spine Surgery" by (L. Jackson et al., 2019) presents a thorough examination of the relationship between preoperative psychological conditions, particularly depression, and postoperative outcomes in spine surgery patients. Through systematic reviews of relevant literature from the PubMed database and the Cochrane Library, the authors sought to clarify whether preoperative treatment of psychological comorbidities could positively influence surgical outcomes.

The findings discussed in this article reveal a significant correlation between psychological disorders and adverse postoperative outcomes, including higher rates of postoperative delirium, increased readmission rates, prolonged hospital stays, and nonroutine hospital discharges. Notably, the authors highlight the specific impact of depression on postoperative delirium,

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

suggesting that individuals with depression may exhibit elevated cortisol levels, which could contribute to this complication ((L. Jackson et al., 2019)). This physiological response underscores the importance of addressing mental health prior to surgical interventions, as it may play a critical role in recovery trajectories. Moreover, the article emphasizes the economic implications of these findings, noting that the increased rates of delirium and readmissions among patients with psychological disorders can lead to higher healthcare costs. This aspect of the study is particularly relevant in the current healthcare environment, where cost-effectiveness is a key consideration in treatment planning. By illustrating the potential financial burden associated with neglecting preoperative psychological health, the authors make a compelling case for the integration of mental health assessments into preoperative evaluations.

However, while the article effectively outlines the associations between psychological comorbidities and surgical outcomes, it would benefit from a more nuanced exploration of the mechanisms underlying these relationships. For instance, further investigation into how specific types of psychological disorders, beyond depression, may influence surgical outcomes could enhance the understanding of this complex interplay. Additionally, the article could expand on the implications of these findings for clinical practice, particularly regarding the development of targeted interventions for patients with identified psychological comorbidities.

The article "Influence of preoperative depression on clinical outcomes after cervical laminoplasty: A retrospective study" by (Zhang et al., 2023) provides essential insights into the impact of preoperative depression on surgical outcomes, particularly in the context of cervical laminoplasty. The authors emphasize that depression is a highly prevalent mental disorder, and their findings suggest a significant correlation between preoperative depression and postoperative recovery.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the critical contributions of this study is the retrospective analysis of clinical outcomes in patients undergoing cervical laminoplasty, which serves as an important reference point for understanding similar implications in lumbar spine surgery. The authors report that patients with preoperative depression exhibited worse postoperative improvement, which aligns with existing literature on the overall influence of mental health on surgical recovery. This finding is particularly relevant for practitioners, as it underscores the necessity of screening for depression prior to surgical interventions.

The methodology employed in the study, while robust, raises questions about the generalizability of the findings to lumbar spine surgery specifically. The focus on cervical laminoplasty may limit the applicability of the results to other types of spinal surgeries, including lumbar fusion, which presents unique challenges and recovery trajectories. Nevertheless, the study effectively highlights the potential adverse effects of untreated depression on surgical outcomes, suggesting that mental health assessments should be integral to preoperative evaluations.

Furthermore, the implications of these findings extend beyond clinical practice to inform future research directions. The authors advocate for further investigations into the mechanisms by which preoperative depression affects surgical outcomes, as well as the development of targeted interventions to address mental health prior to surgery. This perspective is crucial, as it opens avenues for enhancing patient care through a more holistic approach that considers both physical and psychological health.

CONSCLUSION

The literature reviewed emphasizes the significant impact of preoperative depression on surgical outcomes for patients undergoing lumbar spine surgery. The studies collectively highlight a consistent link between depressive symptoms and adverse postoperative results, suggesting that psychological health is a critical factor influencing recovery trajectories.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The foundational study by (Sinikallio et al., 2010) provides compelling evidence that both preoperative and postoperative depressive symptoms predict poor surgical outcomes. This research underscores the importance of mental health not only before surgery but also during the rehabilitation process, indicating that ongoing depressive symptoms can exacerbate recovery challenges. The longitudinal design of this study enhances the credibility of its findings, advocating for continuous mental health monitoring throughout the surgical journey.

Further expanding on this theme, (L. Jackson et al., 2019) conducted a systematic review that connects psychological comorbidities with postoperative complications. Their work reveals that preoperative depression correlates with increased rates of delirium, readmissions, and prolonged hospital stays, suggesting that elevated cortisol levels in depressed individuals may be a contributing factor to these adverse outcomes. This review not only reinforces the need for psychological assessments prior to surgery but also highlights the economic implications of neglecting mental health, as it can lead to higher healthcare costs due to complications.

Additionally, the retrospective study by (Zhang et al., 2023) specifically examines the effects of preoperative depression on outcomes following cervical laminoplasty, providing valuable insights that can be extrapolated to lumbar spine surgeries. The findings indicate that patients with preoperative depression experience significantly worse postoperative improvement, further affirming the necessity of mental health screenings prior to surgical interventions.

In conclusion, the literature consistently demonstrates that preoperative depression is a significant predictor of surgical outcomes in lumbar spine surgery. The integration of psychological assessments into preoperative evaluations is essential for optimizing patient care and enhancing recovery trajectories. Addressing mental health prior to surgery could mitigate adverse outcomes and improve overall patient well-being,

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

highlighting the need for targeted interventions in this population.

DISCLOSURE STATEMENT

Disclosure Statement : The authors have no conflicts of Interest to declare.

REFERENCES

1. Sinikallio, S., M Lehto, S., Aalto, T., Airaksinen, O., Kröger, H., & Viinamäki, H., 2010. Depressive symptoms during rehabilitation period predict poor outcome of lumbar spinal stenosis surgery: A two-year perspective. ncbi.nlm.nih.gov
2. L. Jackson, K., Rumley, J., Griffith, M., Agochukwu, U., & DeVine, J., 2019. Correlating Psychological Comorbidities and Outcomes After Spine Surgery. ncbi.nlm.nih.gov
3. Zhang, W., Sun, T., Wang, S., Zhang, J., Yang, M., & Li, Z., 2023. Influence of preoperative depression on clinical outcomes after cervical laminoplasty: A retrospective study. ncbi.nlm.nih.gov

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved