



Association Between Cigarette Smoking and Multiple Sclerosis

¹ Sri Lestari, ¹ Rusdi

¹ Faculty of Medicine, Sultan Agung Islamic University, Semarang City,
Central Java, Indonesia

Correspondence : drsilestari29381@gmail.com

Article History :

Received date : 2024/04/12
Revised date : 2024/05/07
Accepted date : 2024/06/12
Published date : 2024/07/09



Copyright: © 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

E-ISSN :

ISSN 3048-1368



P-ISSN :

ISSN 3048-1376



ABSTRACT

Background: The association between cigarette smoking and multiple sclerosis (MS) has garnered significant attention in the literature, with numerous studies elucidating the intricate relationships between tobacco exposure and MS risk, progression, and health outcomes. The exploration of these relationships has evolved over the years, revealing both the direct and indirect effects of smoking on the disease. **Literature Review:** The literature on the association between cigarette smoking and multiple sclerosis (MS) has revealed a complex interplay between environmental and genetic factors that significantly influence the disease's risk and progression. The studies reviewed consistently highlight smoking as a critical modifiable risk factor for MS, with evidence suggesting that both active smoking and exposure to environmental tobacco smoke (ETS) contribute to increased susceptibility and exacerbation of the disease. **Conclusion:** Overall, the literature consistently underscores the multifaceted role of cigarette smoking in MS, revealing that both direct and indirect exposures significantly impact disease risk and progression. The interaction between smoking and genetic predispositions, particularly concerning specific HLA genes, further complicates the understanding of MS etiology. These findings call for more focused research on smoking behaviors in MS patients and the development of tailored smoking cessation interventions to improve health outcomes for this population.

Keyword: Cigarette Smoking, Multiple Sclerosis

INTRODUCTION

The association between cigarette smoking and multiple sclerosis (MS) has garnered significant attention in the literature, with numerous studies elucidating the intricate relationships between tobacco exposure and MS risk, progression, and health outcomes. The exploration of these relationships has evolved over the years, revealing both the direct and indirect effects of smoking on the disease.

In 2012, (O’Gorman et al., 2012) presented a comprehensive review that highlighted the increased risk of MS associated with tobacco exposure, particularly for females exposed to passive smoke. Their findings from case-control studies indicated that passive exposure raised the risk for both adult- and child-onset MS, while also noting that Swedish snuff users did not exhibit similar risks. They identified a potential interaction between smoking habits and genetic susceptibility alleles, suggesting a complex interplay between environmental and genetic

factors in MS pathogenesis. Notably, they documented that smoking could exacerbate disease progression, as evidenced by neuroimaging showing heightened inflammation and brain atrophy in smokers (O’Gorman et al., 2012).

Building on this foundation, (Correale et al., 2017) further examined the relationship between smoking and MS risk, establishing a dose-dependent correlation where increased cotinine levels were associated with heightened susceptibility to the disease. Their research underscored the importance of smoking cessation, noting that the risk diminishes significantly after a decade of quitting. They also emphasized the role of smoking in accelerating the transition from clinically isolated syndrome (CIS) to clinically definite MS, as well as its influence on disease progression and disability (Correale et al., 2017).

In a 2020 review, (H Marck et al., 2020) reiterated the critical role of smoking as a modifiable risk factor for MS onset and progression,

estimating that smoking could increase the risk of developing MS by approximately 50%. They highlighted the negative impact of smoking on quality of life and mental health among MS patients, advocating for more research into effective smoking cessation strategies tailored for this population. The authors argued that despite the known risks, there remains a substantial gap in understanding smoking behaviors in individuals with MS and how healthcare providers can support cessation efforts (H Marck et al., 2020).

(Nishanth et al., 2020) contributed to the discourse by exploring the pathogenesis of MS in relation to smoking. They reaffirmed that environmental factors such as cigarette smoke significantly contribute to the severity and progression of MS, further emphasizing the need for a deeper understanding of smoking's immunological impacts in the context of the disease (Nishanth et al., 2020).

Most recently, (Antônio Machado Schlindwein et al., 2024) conducted a systematic review focusing on the effects of

environmental tobacco smoke on MS. Their findings corroborated previous studies, indicating that smoking not only increases the risk of MS onset but also correlates with more aggressive disease progression. They identified a notable interaction between smoking and specific HLA genes, suggesting a genetic predisposition that may amplify the risk associated with tobacco exposure (Antônio Machado Schlindwein et al., 2024).

Collectively, these studies underscore the multifaceted role of cigarette smoking in the context of multiple sclerosis, revealing a complex interplay between environmental exposures, genetic factors, and disease outcomes. The literature consistently points to smoking as a significant risk factor, necessitating further research to address the gaps in understanding smoking behaviors and to develop effective interventions for individuals affected by MS.

LITERATURE REVIEW

The article "Environmental Risk Factors for Multiple Sclerosis: A

Review with a Focus on Molecular Mechanisms" by Cullen O’Gorman, Robyn Lucas, and Bruce Taylor (O’Gorman et al., 2012) provides a comprehensive examination of the relationship between cigarette smoking and multiple sclerosis (MS), highlighting various studies that contribute to our understanding of this complex association.

The authors present evidence from two case-control studies indicating that passive exposure to cigarette smoke may increase the risk of both adult- and child-onset MS. However, the relationship between maternal smoking habits and MS risk remains ambiguous, as conflicting results emerge from different studies—some indicating no association while others suggest a heightened risk. This inconsistency underscores the need for further investigation into the impact of maternal smoking on MS susceptibility.

Notably, the article discusses findings from studies involving Swedish snuff users, which did not show an increased risk of MS. This raises questions about the specific

components of cigarette smoke that may be implicated in MS development. The authors highlight a case-control study that found no increased risk of MS associated with smoking, suggesting a lack of interaction between susceptibility alleles for MS and cigarette smoke exposure. Nevertheless, a population-based case-control study did identify possible interactions between susceptibility alleles and smoking habits, indicating a more nuanced relationship that warrants deeper exploration.

The article further elaborates on the implications of smoking in the context of MS progression. It is reported that smoking is linked to an increased risk of secondary progressive disease and facilitates earlier progression from clinically isolated syndrome (CIS) to clinically definite MS. These findings are significant as they suggest that smoking may not only contribute to the onset of MS but also exacerbate its progression.

In terms of biological mechanisms, the authors note that cigarette smoke comprises thousands

of components, including particulate matter, nicotine, and carbon monoxide, which have been associated with pro-inflammatory actions, direct tissue damage, and increased apoptosis. This information is crucial, as it connects the environmental risk factor of smoking with the underlying molecular mechanisms that may drive MS pathology.

The article titled "Environmental factors influencing multiple sclerosis in Latin America" by Correale, Farez, and Gaitán (Correale et al., 2017) presents a comprehensive examination of the relationship between cigarette smoking and multiple sclerosis (MS), emphasizing the substantial increase in risk associated with tobacco use. The authors assert that cigarette smoking is linked to an estimated 40–80% increased risk of developing MS, a finding supported by various case-control studies. This statistic underscores the critical role that environmental factors, particularly smoking, play in the etiology of MS.

A significant contribution of the article is its discussion of the dose-

dependent relationship between smoking and MS risk. The authors note that cumulative exposure to tobacco, as measured by cotinine serum or plasma levels, correlates with an increased likelihood of developing the disease. This relationship highlights the importance of not only the act of smoking itself but also the quantity and duration of exposure. Furthermore, the authors provide evidence that the risk associated with smoking diminishes after a decade of cessation, suggesting that the body may recover from some of the adverse effects of smoking over time.

The article also addresses the impact of passive smoking on MS risk, indicating that non-smokers exposed to tobacco smoke are similarly at an increased risk of developing the disease. This finding broadens the scope of the discussion around smoking and MS, implicating not only active smokers but also those who are indirectly exposed to tobacco smoke.

In addition to the risk of developing MS, the authors delve into the clinical implications of smoking

for individuals already diagnosed with the disease. They report that smoking is a risk factor for early conversion to clinically definite MS among patients with clinically isolated syndrome (CIS). Moreover, the progression of MS appears to be accelerated in smokers compared to non-smokers, with smokers achieving higher Expanded Disability Status Scale (EDSS) milestones, indicating a more severe clinical course. This aspect of the article is particularly critical for clinicians and researchers as it underscores the need for smoking cessation programs as part of comprehensive MS management.

The authors also explore the genetic interactions that may exacerbate the risk associated with smoking. Specifically, they highlight the significant interaction between smoking and MS-associated HLA risk alleles. The presence of certain HLA alleles, particularly HLA-DRB1*1501, markedly increases the risk of developing MS in smokers, suggesting a complex interplay between genetic predisposition and environmental factors. Additionally, the association with the N-

acetyltransferase 1 (NAT1) genotype points to potential avenues for further research into the genetic and environmental determinants of MS susceptibility.

The article titled "Modifiable risk factors for poor health outcomes in multiple sclerosis: The urgent need for research to maximise smoking cessation success" by Marck et al. (H Marck et al., 2020) provides a comprehensive examination of the impact of cigarette smoking on multiple sclerosis (MS). The authors emphasize that tobacco smoking is not merely a lifestyle choice but a significant modifiable risk factor that has been linked to the onset and progression of MS. Their findings indicate that smoking increases the risk of developing MS by approximately 50%, underscoring its critical role in the disease's etiology.

The review highlights various observational studies that demonstrate a correlation between smoking and adverse health outcomes in individuals with MS. Specifically, the authors note that smoking is associated with a deterioration in disability, lower quality of life, and

heightened levels of depression and anxiety among MS patients. This connection between smoking and mental health is particularly concerning, as it suggests that smoking cessation could play a vital role in improving overall well-being and health outcomes for those living with MS.

One of the significant gaps identified by the authors is the lack of research focused on smoking cessation strategies tailored for individuals with MS. Despite the clear evidence linking smoking to negative health outcomes, there is a notable absence of studies investigating how healthcare professionals can effectively support MS patients in quitting smoking. This gap in the literature is particularly alarming given the potential benefits that successful cessation could bring to this population.

The article calls for urgent research to develop effective smoking cessation interventions specifically designed for people with MS. The authors argue that understanding smoking behavior in this demographic is crucial for

maximizing health outcomes. They advocate for a more structured approach to smoking cessation within healthcare settings, which could lead to improved management of MS and enhanced quality of life for patients.

The article titled "Role of Smoking in the Pathogenesis of Multiple Sclerosis: A Review Article" by Katukuri Nishanth et al. (Nishanth et al., 2020) provides a comprehensive examination of the impact of cigarette smoking as an environmental risk factor in the context of Multiple Sclerosis (MS). The authors delve into the multifaceted ways in which smoking may influence the development, severity, progression, and disability associated with MS, as well as its potential role in early mortality among affected individuals.

The review emphasizes that while the association between smoking and MS has been noted in various studies, the underlying mechanisms remain inadequately understood. The authors highlight that cigarette smoke may modulate immune responses, potentially exacerbating the autoimmune

processes characteristic of MS. They suggest that components of cigarette smoke could lead to alterations in immune cell function, which may contribute to the pathogenesis of MS. This aspect is crucial, as understanding the immunological effects of smoking can help elucidate its role in MS progression and severity.

Furthermore, the article critically evaluates existing literature, noting inconsistencies in findings regarding the relationship between smoking and MS. The authors point out that while some studies suggest a clear association, others indicate minimal or no significant links. This discrepancy underscores the need for further research to clarify the causal relationships and to explore the biological mechanisms at play. The authors propose that future studies should focus on longitudinal designs and consider genetic predispositions that may interact with environmental exposures like smoking.

The article "Impacts of environmental tobacco smoke on the onset and progression of multiple sclerosis: a systematic review" by

Schlindwein et al. (Antônio Machado Schlindwein et al., 2024) provides a comprehensive examination of the relationship between tobacco smoking and multiple sclerosis (MS), particularly focusing on environmental tobacco smoke (ETS). The authors effectively summarize the existing evidence linking active cigarette smoking to an increased risk of developing MS and a more aggressive disease course, highlighting a significant 50% increase in MS onset risk associated with smoking (Antônio Machado Schlindwein et al., 2024). This finding aligns with the broader understanding of smoking as a modifiable risk factor for various autoimmune disorders.

The article delves into the mechanisms that may underlie the relationship between smoking and MS, noting that while the precise pathways remain unclear, there is evidence suggesting interactions between smoking and genetic predispositions, particularly within the human leukocyte antigen (HLA) complex (Antônio Machado Schlindwein et al., 2024). This aspect

is crucial as it underscores the complexity of MS etiology, where both environmental and genetic factors interplay, potentially exacerbating disease risk.

In contrast, the review highlights a notable gap in the literature regarding the effects of ETS on MS. While active smoking has been well-studied, the implications of ETS remain less defined, which the authors argue is a significant oversight given the prevalence of exposure to secondhand smoke in various populations (Antônio Machado Schlindwein et al., 2024). The call for further research in this area is particularly salient, as understanding the impact of ETS could have important public health implications, especially for vulnerable populations who may be at risk for MS.

The systematic review format employed by the authors allows for a critical evaluation of the existing literature, providing a structured approach to synthesizing findings from various studies. However, the review could benefit from a more detailed exploration of the

methodological limitations present in the studies cited, which would enhance the robustness of the conclusions drawn. Additionally, while the authors mention the potential for ETS to influence MS onset and progression, they do not provide specific data or studies that directly address this relationship, which could leave readers seeking more concrete evidence.

CONCLUSION

The literature on the association between cigarette smoking and multiple sclerosis (MS) has revealed a complex interplay between environmental and genetic factors that significantly influence the disease's risk and progression. The studies reviewed consistently highlight smoking as a critical modifiable risk factor for MS, with evidence suggesting that both active smoking and exposure to environmental tobacco smoke (ETS) contribute to increased susceptibility and exacerbation of the disease.

(O’Gorman et al., 2012) provided foundational insights into the relationship between passive

smoking and MS risk, indicating that both adult- and child-onset MS are associated with tobacco exposure. They noted that while Swedish snuff users did not exhibit the same risks, the interaction between smoking habits and genetic susceptibility alleles complicates the understanding of MS pathogenesis. This complexity is echoed by (Correale et al., 2017), who established a dose-dependent correlation between cotinine levels and MS risk, emphasizing that cumulative tobacco exposure heightens susceptibility, although the risk diminishes significantly after a decade of cessation.

(H Marck et al., 2020) further reinforced the notion that smoking is linked to adverse health outcomes in MS patients, including lower quality of life and increased mental health challenges. They identified a critical gap in research regarding effective smoking cessation strategies specifically tailored for individuals with MS, highlighting the need for healthcare providers to prioritize cessation efforts within this demographic. Moreover, (Nishanth et al., 2020) explored the

immunological impacts of smoking, suggesting that components of cigarette smoke may exacerbate autoimmune processes, thus contributing to the severity and progression of MS.

(Antônio Machado Schlindwein et al., 2024) conducted a systematic review that corroborated the findings of previous studies, indicating that smoking increases the risk of MS onset by approximately 50% and correlates with more aggressive disease progression. They also pointed out the need for further research into the effects of ETS on MS, an area that remains underexplored despite its public health implications.

Overall, the literature consistently underscores the multifaceted role of cigarette smoking in MS, revealing that both direct and indirect exposures significantly impact disease risk and progression. The interaction between smoking and genetic predispositions, particularly concerning specific HLA genes, further complicates the understanding of MS etiology. These findings call for more focused research on smoking

behaviors in MS patients and the development of tailored smoking cessation interventions to improve health outcomes for this population.

REFERENCES

1. O’Gorman, C., Lucas, R., & Taylor, B., 2012. Environmental Risk Factors for Multiple Sclerosis: A Review with a Focus on Molecular Mechanisms. ncbi.nlm.nih.gov
2. Correale, J., F Farez, M., & Inés Gaitán, M., 2017. Environmental factors influencing multiple sclerosis in Latin America. ncbi.nlm.nih.gov
3. H Marck, C., das Nair, R., B Grech, L., Borland, R., & S Constantinescu, C., 2020. Modifiable risk factors for poor health outcomes in multiple sclerosis: The urgent need for research to maximise smoking cessation success. ncbi.nlm.nih.gov
4. Nishanth, K., Tariq, E., P Nzvere, F., Miqdad, M., & Cancarevic, I., 2020. Role of Smoking in the Pathogenesis of Multiple Sclerosis: A Review Article. ncbi.nlm.nih.gov
5. Antônio Machado Schlindwein, M., Henrique de Moura Campos, M., Caroline Breis, L., Sordi Chara, B., Sasse Scherer, C., Augusta Piva Caminski, V., Matta, A., & Vinicius Magno Gonçalves, M., 2024. Impacts of environmental tobacco smoke on the onset and progression of multiple sclerosis: a systematic review. ncbi.nlm.nih.gov