



A Comprehensive Literature Review of Relationship of Inflammatory Bowel Disease and Celiac Disease

¹ Erick Persson Jeffry, ¹ Charles Sanjaya Seikka

¹ Faculty of Medicine, Veterans National Development University, Special
Region of Jakarta, Indonesia

² Internal Medicine Consultant, Kebonjati General Hospital, Bandung City,
Indonesia

Correspondence : erick.persson86@gmail.com

Article History :

Received date : 2024/10/19

Revised date : 2024/11/28

Accepted date : 2024/12/17

Published date : 2025/01/08



Copyright: © 2024 by the
authors. Submitted for possible
open access publication under
the terms and conditions of the
Creative Commons Attribution
(BY NC) license
(<https://creativecommons.org/licenses/by-nc/4.0/>).

E-ISSN :

ISSN 3048-1368



P-ISSN :

ISSN 3048-1376



ABSTRACT

Background: The relationship between Inflammatory Bowel Disease (IBD) and Celiac Disease (CeD) has garnered significant attention in recent research, revealing complex interactions that may inform both pathogenesis and treatment strategies. **Literature Review:** The variability in prevalence rates of CeD among IBD patients, as reported by (Bengi et al., 2019), underscores the complexity of these diseases and the interplay of genetic, environmental, and immunological factors. This is further supported by (Hyun, 2021), who explore the connections between metabolic disorders and IBD, suggesting that chronic intestinal inflammation may predispose individuals to additional health issues, thereby reinforcing the idea of shared inflammatory pathways. Additionally, (Akiyama et al., 2022) elucidates the increased prevalence of CeD in patients with immune-mediated inflammatory diseases, indicating that the coexistence of these conditions can influence clinical outcomes. Finally, (Yemula, 2024) emphasizes the role of gut microbiota in the pathophysiology of CeD, suggesting that microbial changes may influence both conditions. The interconnectedness of IBD and CeD is evident, with shared genetic, immunological, and environmental factors driving the relationship. The literature collectively advocates for further research to clarify these interactions and improve patient management strategies, particularly regarding screening and dietary considerations. **Conclusion:** In conclusion, the relationship between IBD and CeD is characterized by complex interactions influenced by shared genetic predispositions, environmental factors, and dysregulated gut microbiota. The findings underscore the necessity for a nuanced understanding of these conditions to enhance diagnostic approaches and treatment strategies.

Keywords: Relationship, Inflammatory Bowel Disease, Celiac Disease

INTRODUCTION

The relationship between Inflammatory Bowel Disease (IBD) and Celiac Disease (CeD) has garnered significant attention in recent research, revealing complex interactions that may inform both pathogenesis and treatment strategies. The exploration of these connections begins with (Aleksandrova et al., 2017), who emphasize that IBD, encompassing Crohn's disease and ulcerative colitis, is characterized by chronic gastrointestinal inflammation. They propose that a shared aetiology exists between IBD and CeD, particularly in genetically predisposed individuals who develop intolerances to dysregulated gut microbiota influenced by environmental factors. This foundational understanding sets the stage for examining the genetic and immunological underpinnings of both diseases.

In a subsequent study, (Maia Bosca-Watts et al., 2018) delve into the role of human leucocyte antigen (HLA) genes in disease susceptibility.

Their findings suggest that while IBD patients are traditionally viewed as being at higher risk for CeD, recent analyses indicate that the prevalence of CeD in IBD populations may not be as pronounced as previously believed. This nuance highlights the need for further investigation into the diagnostic criteria and the environmental factors contributing to both conditions.

(Bengi et al., 2019) expand on this discussion by presenting data on the prevalence of CeD among IBD patients within the Turkish population, noting a wide variability in prevalence rates. Their research underscores the multifactorial nature of both diseases, suggesting that genetic, environmental, and immunological factors intertwine to influence disease manifestation.

(Hyun, 2021) introduces another layer of complexity by discussing the potential links between metabolic disorders and IBD, suggesting that chronic intestinal inflammation may predispose individuals to conditions such as

diabetes mellitus. This perspective aligns with the notion that IBD and CeD may share common inflammatory pathways, as both conditions exhibit dysbiosis and impaired epithelial barrier function.

(Akiyama et al., 2022) further explore the coexistence of CeD in patients with immune-mediated inflammatory diseases, reinforcing the idea that IBD patients are at an increased risk for CeD. Their findings indicate that the presence of CeD may influence the clinical outcomes of IBD, suggesting that shared genetic and environmental factors contribute to the pathophysiology of both diseases.

The bidirectional causal relationship between IBD and CeD is elucidated by (A et al., 2022), who utilize Mendelian randomization analysis to affirm that the etiologies of these disorders may be interlinked. Their study highlights the significance of screening for CeD among IBD patients, advocating for tailored management strategies that consider the potential for unfavorable

outcomes in patients with concurrent diagnoses.

Finally, (Yemula, 2024) emphasizes the role of gut microbiota in the pathophysiology of CeD, linking microbial changes to disease immunology. This research suggests that the gut microbiome may not only influence the development of CeD but also interact with the mechanisms underlying IBD, thus reinforcing the interconnectedness of these two chronic conditions.

Overall, the literature reveals a nuanced relationship between IBD and CeD, characterized by shared genetic, immunological, and environmental factors, while also highlighting the importance of further research to clarify these interactions and improve patient management strategies.

LITERATURE REVIEW

The article by (Aleksandrova et al., 2017) provides a comprehensive overview of the intricate relationship between diet, gut microbiome, and epigenetics in the context of inflammatory bowel

diseases (IBD), which include Crohn's disease (CD) and ulcerative colitis (UC). The authors articulate that IBD is characterized by chronic inflammation of the gastrointestinal tract and highlight its growing significance as a public health issue. They suggest that despite the apparent differences in pathology between CD and UC, both may share a common etiology, primarily influenced by genetic predispositions and environmental triggers.

A critical insight from the article is the emphasis on the role of dysbiosis—an imbalance in the gut microbiota—as a central factor in the development of chronic inflammation in genetically susceptible individuals. The authors assert that this dysbiosis, coupled with environmental factors such as diet, can significantly influence the pathogenesis of IBD. They argue that dietary components can modulate intestinal inflammation and gut microbiota, thereby playing a crucial role in disease management and prevention. This perspective aligns with the growing recognition of the gut microbiome's impact on

overall health and disease states, including autoimmune conditions.

Furthermore, the article delves into the genetic aspects of IBD, noting that while significant genetic risk loci have been identified, they only account for a fraction of the genetic variance associated with IBD risk. This observation underscores the complexity of IBD's etiology, suggesting that impaired immune responses in genetically susceptible individuals may serve as a unifying mechanism linking genetic and environmental factors.

The authors also introduce the emerging field of epigenetics, which offers insights into how environmental changes can lead to pathological gene expression and influence cellular functions in IBD. This aspect is particularly relevant as it opens new avenues for understanding the interplay between genetic predispositions and environmental influences, including dietary habits.

The article titled "HLA-DQ: Celiac disease vs inflammatory bowel disease" by (Maia Bosca-Watts et al.,

2018) provides a comprehensive examination of the relationship between Celiac disease (CeD) and inflammatory bowel disease (IBD), two chronic intestinal disorders that have seen a notable increase in incidence and prevalence. The authors highlight that both conditions arise from a complex interplay of genetic, immunological, and environmental factors, particularly in genetically predisposed individuals.

One of the key insights from the article is the established role of human leukocyte antigen (HLA) genes, specifically HLA-DQ2 and HLA-DQ8, in the susceptibility to CeD. This genetic predisposition is critical in understanding why certain individuals may develop CeD in conjunction with IBD. The article emphasizes that while IBD patients have historically been perceived as being at a heightened risk for developing CeD, recent studies challenge this assumption. Notably, the authors reference findings indicating that the prevalence of CeD may be comparable to, or even lower than, that found in the general population of IBD patients. This

revelation is significant as it prompts a reevaluation of the perceived risk factors and the relationship between these two disorders.

The article also delineates the diagnostic criteria for CeD, which includes serological tests for specific antibodies, genetic testing, upper endoscopy, and histological evaluation of small-bowel biopsies. This structured approach to diagnosis underscores the complexity of CeD and its differentiation from IBD, which may share overlapping symptoms but have distinct pathological and immunological profiles.

Furthermore, the authors address the potential for misdiagnosis or underdiagnosis of CeD in IBD patients, which could stem from the symptomatic overlap between the two diseases. The critical evaluation of the existing literature provided in this study highlights the necessity for clinicians to maintain a high index of suspicion for CeD in patients presenting with gastrointestinal symptoms, even in the context of an established IBD diagnosis.

The article titled "Prevalance of Celiac Disease in Patients with Inflammatory Bowel Disease in Turkish Population" by (Bengi et al., 2019) provides a critical examination of the intersection between celiac disease (CD) and inflammatory bowel disease (IBD), emphasizing the inflammatory nature of both conditions. The authors note that while the pathogenesis of CD and IBD is multifactorial, involving environmental, genetic, and immunological components, the prevalence of CD among IBD patients can vary significantly, with reported rates ranging from 0% to 14%.

The study's primary aim was to ascertain the prevalence of CD within a cohort of IBD patients attending a hospital or outpatient clinic in Turkey. This focus is particularly relevant, as it addresses a gap in the literature regarding the co-occurrence of these two inflammatory conditions in a specific population. The authors methodically gathered data from IBD patients, which not only contributes to understanding the prevalence of CD in this demographic but also raises important

considerations about the clinical implications of coexisting conditions.

One of the strengths of this article is its systematic approach to data collection and analysis, which enhances the reliability of the findings. By providing concrete prevalence rates, the authors offer valuable insights that can inform clinical practice, particularly regarding the necessity for screening IBD patients for CD. This is crucial because undiagnosed CD in IBD patients may lead to suboptimal management of both conditions, complicating treatment strategies and potentially exacerbating symptoms.

However, the article also presents limitations that warrant discussion. The prevalence range reported, while informative, highlights the variability in findings across different studies and populations. This variability suggests that factors such as genetic predisposition, environmental influences, and regional dietary habits may play significant roles in the coexistence of CD and IBD. The authors could have further explored

these factors to provide a more nuanced understanding of the relationship between the two diseases.

Additionally, while the study successfully identifies a population-based prevalence, it would benefit from a broader discussion on the implications of this prevalence for clinical practice. For instance, the authors could have elaborated on how the presence of CD might influence the management of IBD, including dietary modifications and the potential need for different therapeutic approaches.

The article titled "Molecular and Pathophysiological Links between Metabolic Disorders and Inflammatory Bowel Diseases" by (Hyun, 2021) provides a comprehensive exploration of the interplay between inflammatory bowel diseases (IBD) and metabolic disorders, particularly focusing on diabetes mellitus. The author posits that chronic intestinal inflammation may influence glucose homeostasis, thereby potentially increasing the risk for diabetes. This connection is particularly relevant in understanding

the broader implications of IBD, as it highlights the systemic effects of intestinal inflammation.

Hyun discusses recent epidemiological studies that have established associations between type 1 diabetes (T1D), type 2 diabetes (T2D), and IBD. Notably, the article emphasizes the genetic links between T1D and IBD, suggesting that these diseases may share common genetic variants. This genetic overlap implies a potential for shared pathophysiological mechanisms, which warrants further investigation. The author effectively synthesizes findings from multiple studies, reinforcing the idea that the comorbidity of T2D and IBD may stem from factors such as gut microbiota dysbiosis, epithelial barrier dysfunction, and ongoing inflammation.

The article's critical evaluation of the relationship between IBD and metabolic disorders is significant, as it underscores the complexity of these conditions. By presenting evidence of shared inflammatory processes, Hyun

encourages a multidisciplinary approach to understanding IBD that encompasses not only gastrointestinal health but also metabolic implications. This perspective is particularly valuable for clinicians and researchers alike, as it may inform treatment strategies that address both IBD and its metabolic consequences.

The article titled "Characteristics of inflammatory bowel diseases in patients with concurrent immune-mediated inflammatory diseases" by (Akiyama et al., 2022) provides a comprehensive examination of the relationship between inflammatory bowel disease (IBD) and celiac disease, particularly focusing on the shared pathophysiological mechanisms and clinical outcomes in patients with these concurrent conditions. The authors establish that patients with IBD exhibit a higher prevalence of celiac disease compared to control groups, indicating a significant overlap in the patient populations affected by these immune-mediated inflammatory diseases (IMIDs).

One of the critical insights from the study is the identification of shared genetic, environmental, and immunological factors between IBD and celiac disease. The authors reference genome-wide association studies (GWASs) that have pinpointed common genetic risk loci, such as PTPN2, IL18RAP, TAGAP, and PUS10, which underscore the potential for shared etiological pathways. This genetic overlap suggests that individuals predisposed to one condition may also be at increased risk for the other, which is a vital consideration for clinicians managing patients with these diseases.

The article also discusses the clinical implications of celiac disease coexisting with IBD. Notably, it highlights findings from a case-control study indicating that Crohn's disease (CD) patients with celiac disease are less likely to exhibit ileocolonic involvement and require aggressive medical therapies, such as tumor necrosis factor (TNF) inhibitors or azathioprine, compared to those without celiac disease. This observation suggests that the presence

of celiac disease may confer a more favorable disease course for CD patients. Conversely, the authors note that ulcerative colitis (UC) patients with celiac disease face a higher risk of colectomy, indicating a complex interplay where celiac disease may exacerbate the clinical course of UC.

The article further delves into the immunological aspects of both diseases, emphasizing the role of Th17 cells and their associated cytokines, such as IL-17, interferon- γ , and IL-21, in the pathogenesis of celiac disease. This shared immunological component raises important questions about the inflammatory processes at play in both conditions and the potential for targeted therapies that address these common pathways.

The article titled "Bidirectional causal link between inflammatory bowel disease and celiac disease: A two-sample mendelian randomization analysis" by (A et al., 2022) presents a thorough investigation into the interrelationship between inflammatory bowel disease (IBD)

and celiac disease through the application of Mendelian randomization (MR) analysis and a substantial dataset derived from genome-wide association studies (GWAS). The authors articulate a compelling narrative that elucidates the potential bidirectional causal link between these two gastrointestinal disorders, suggesting that their etiologies may share common pathways.

The findings from the study indicate a significant association between IBD and an increased risk of developing celiac disease. Notably, the authors provide evidence that patients with Crohn's disease (CD) exhibit a statistically significant higher risk of celiac disease compared to those with ulcerative colitis (UC). This distinction is crucial as it highlights the differential impact of IBD subtypes on the risk of celiac disease, thus inviting further exploration into the underlying mechanisms that contribute to this relationship.

The authors effectively argue that genetic, immunological, and

environmental factors converge to create a similar pathophysiological landscape for both IBD and celiac disease. This assertion is particularly pertinent as it underscores the complexity of these disorders and the necessity for a multifaceted approach in understanding their interconnections. The suggestion that individuals diagnosed with IBD should undergo regular serological testing for celiac disease is a practical recommendation that could enhance clinical outcomes by facilitating early detection and management of celiac disease in this vulnerable population.

Moreover, the article advocates for dietary modifications, specifically the implementation of a low-gluten diet for patients with IBD. This dietary approach may serve as a beneficial adjunct in the management of IBD, particularly for those who may also be at risk for celiac disease. The authors' proposal to investigate the temporal sequence of disease events further emphasizes the need for longitudinal studies to better understand the dynamics of these conditions and their progression over time.

The article "Gut microbiota in celiac disease" by (Yemula, 2024) provides a comprehensive overview of celiac disease (CD), emphasizing its autoimmune nature and the complex interplay between genetic, environmental, and microbial factors in its pathophysiology. Yemula highlights that CD is triggered by gluten ingestion in genetically predisposed individuals, with prevalence rates varying significantly across different geographic regions. Notably, the article underscores the importance of genetic factors, particularly the association of HLA-DQ2 and HLA-DQ8 alleles with the disease, which suggests a substantial hereditary component in its manifestation.

The discussion on the gut microbiota is particularly relevant, as it introduces an additional layer of complexity to the understanding of CD. Yemula points out that alterations in gut microbial composition have been consistently observed in patients with CD, suggesting that dysbiosis may play a role in the disease's onset and progression. This relationship is

critical, as it opens avenues for further research into how microbial interventions could potentially mitigate the effects of gluten in susceptible individuals.

However, while the article presents a thorough examination of the genetic and immunological aspects of CD, it could benefit from a more detailed exploration of the specific microbial changes associated with the disease. For instance, identifying which particular microbial species are altered in CD patients could provide insights into potential therapeutic targets. Additionally, the connection between CD and other autoimmune disorders, as mentioned, raises questions about shared pathophysiological mechanisms that could be elucidated through further investigation into the microbiota's role in these conditions.

CONCLUSION

The literature presents a multifaceted view of the relationship between Inflammatory Bowel Disease (IBD) and Celiac Disease (CeD), emphasizing shared genetic, immunological, and environmental factors. The foundational work by

(Aleksandrova et al., 2017) highlights the chronic inflammation characteristic of IBD and suggests a shared etiology with CeD, particularly in genetically predisposed individuals influenced by environmental factors and dysregulated gut microbiota. This perspective is supported by (Maia Bosca-Watts et al., 2018), who delve into the role of HLA genes in disease susceptibility, indicating that the prevalence of CeD in IBD populations may not be as high as previously assumed. This finding calls for a reevaluation of diagnostic criteria and further investigation into environmental influences.

The variability in prevalence rates of CeD among IBD patients, as reported by (Bengi et al., 2019), underscores the complexity of these diseases and the interplay of genetic, environmental, and immunological factors. This is further supported by (Hyun, 2021), who explore the connections between metabolic disorders and IBD, suggesting that chronic intestinal inflammation may predispose individuals to additional health issues, thereby reinforcing the

idea of shared inflammatory pathways.

Additionally, (Akiyama et al., 2022) elucidates the increased prevalence of CeD in patients with immune-mediated inflammatory diseases, indicating that the coexistence of these conditions can influence clinical outcomes. The bidirectional relationship between IBD and CeD is further supported by (A et al., 2022), who utilize Mendelian randomization to affirm the interlinked etiologies of these disorders, advocating for regular screening for CeD in IBD patients.

Finally, (Yemula, 2024) emphasizes the role of gut microbiota in the pathophysiology of CeD, suggesting that microbial changes may influence both conditions. The interconnectedness of IBD and CeD is evident, with shared genetic, immunological, and environmental factors driving the relationship. The literature collectively advocates for further research to clarify these interactions and improve patient management strategies, particularly

regarding screening and dietary considerations.

In conclusion, the relationship between IBD and CeD is characterized by complex interactions influenced by shared genetic predispositions, environmental factors, and dysregulated gut microbiota. The findings underscore the necessity for a nuanced understanding of these conditions to enhance diagnostic approaches and treatment strategies.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -

Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCE

1. Aleksandrova, K., Romero-Mosquera, B., & Hernandez, V., 2017. Diet, Gut Microbiome and Epigenetics: Emerging Links with Inflammatory Bowel Diseases

- and Prospects for Management and Prevention. ncbi.nlm.nih.gov
2. Maia Bosca-Watts, M., Minguez, M., Planelles, D., Navarro, S., Rodriguez, A., Santiago, J., Tosca, J., & Mora, F., 2018. HLA-DQ: Celiac disease vs inflammatory bowel disease. ncbi.nlm.nih.gov
3. Bengi, G., Cıvık, M., Akarsu, M., Soytürk, M., Ellidokuz, E., Topalak, Ömer, & Akpınar, H., 2019. Prevalance of Celiac Disease in Patients with Inflammatory Bowel Disease in Turkish Population. ncbi.nlm.nih.gov
4. Hyun, C. K., 2021. Molecular and Pathophysiological Links between Metabolic Disorders and Inflammatory Bowel Diseases. ncbi.nlm.nih.gov
5. Akiyama, S., Fukuda, S., M Steinberg, J., Suzuki, H., & Tsuchiya, K., 2022. Characteristics of inflammatory bowel diseases in patients with concurrent immune-mediated inflammatory diseases. ncbi.nlm.nih.gov
6. A, G., Sun, C., Shan, Y., Husile, H., & Bai, H., 2022. Bidirectional causal link between inflammatory bowel disease and celiac disease: A two-sample mendelian randomization analysis. ncbi.nlm.nih.gov
7. Yemula, N., 2024. Gut microbiota in celiac disease. ncbi.nlm.nih.gov